


1.  $P_{\text{vap}}^{\circ}(\text{H}_2\text{O}) = 0.0317 \text{ bar}$ . Circle all the conditions that will cause water to condense spontaneously.
- Partial pressure of water = 1 bar
  - Partial pressure of water = 0.1 bar
  - Partial pressure of water = 0.01 bar
  - Partial pressure of water = 0.0317 bar, temperature increases
  - Partial pressure of water = 0.0317 bar, volume decreases
2. The vapor pressure of a liquid is 0.92 atm at 60 °C. Circle all the temperatures that could correspond to the boiling point of the liquid.
- 35 °C    45 °C    55 °C    60 °C    65 °C
3. When acetic acid ( $\text{CH}_3\text{COOH}$ ) dissolves in benzene, it forms dimers. A solution of acetic acid in benzene has a freezing point of 5.40 °C. What is the molality of the acetic acid/benzene solution?

$$T_{f,\text{benzene}} = 5.53 \text{ }^{\circ}\text{C} \quad k_{f,\text{benzene}} = 5.12 \frac{\text{ }^{\circ}\text{C}\cdot\text{kg}}{\text{mol}}$$



4. Consider two gases, iodine chloride ( $\text{ICl}(\text{g})$ ) and iodine bromide ( $\text{IBr}(\text{g})$ )
- (a) Circle the one with greater  $k_H$  in water at 25 °C?


$k_H(\text{ICl})$      $k_H(\text{IBr})$

Two separate systems are prepared at the same initial temperature and pressure:

- System 1:  $\text{ICl}(\text{g})$  at 2 bar in 250 mL of water
  - System 2:  $\text{IBr}(\text{g})$  at 2 bar in 250 mL of water
- (b) Suppose you want the systems to have equal concentrations of  $\text{ICl}(\text{aq})$  and  $\text{IBr}(\text{aq})$ . Which system would you compress to achieve this goal?
- $\text{ICl}$      $\text{IBr}$     Cannot be determined
- (c) Suppose you want the systems to have equal concentrations of  $\text{ICl}(\text{aq})$  and  $\text{IBr}(\text{aq})$ . Which system would you heat at constant pressure to achieve this goal?
- $\text{ICl}$      $\text{IBr}$     Cannot be determined

5. Two aqueous solutions are placed in a U-tube separated by a semipermeable membrane. One solution is 1.00 L of 0.05 M urea (a molecular solute). The other is 1.00 L of 0.02 M  $\text{Na}_2\text{SO}_4$ . Assume ideal solutions at the same temperature.
- (a) Which solution has the higher osmotic pressure?
- (b) In order to stop water from flowing, pressure must be applied to which side?
- (c) Water is allowed to flow through the semipermeable membrane until the osmotic pressure of both solutions equalize. What is the concentration of  $\text{Na}_2\text{SO}_4$  when this occurs?

6. Two beakers are placed together in a sealed container. One beaker contains 1.00 L 0.01 m  $\text{NaNO}_3(\text{aq})$ , the other contains 1.00 L 0.01 m  $\text{Fe}(\text{NO}_3)_3(\text{aq})$ . After some time, the system reaches equilibrium as the water vapor causes redistribution between the beakers. Sketch the water levels below.



**Homework Problem 29**

1. The standard (1 bar) boiling point of ethanol ( $\text{C}_2\text{H}_6\text{O}$ ) is  $78.4^\circ\text{C}$ . 9.15 g of a molecular solute is dissolved in 100.0 g of ethanol. The vapor pressure of the resulting solution is 0.97 at  $78.4^\circ\text{C}$ .
- (a) What is the mol fraction of ethanol in the solution?
- (b) What is the molar mass of the solute?
- (c) What is the boiling point of the solution? ( $k_b$  for ethanol =  $1.07 \frac{\text{K}\cdot\text{kg}}{\text{mol}}$ )